

γ -Ray Spectroscopy in Λ Hypernuclei

H. Tamura,^{*1} S. Ajimura,² H. Akikawa,^{3,†} D.E. Alburger,⁴ K. Aoki,⁵ A. Banu,⁶
 R.E. Chrien,⁴ G.B. Franklin,⁷ J. Franz,⁸ Y. Fujii,¹ Y. Fukao,³ T. Fukuda,⁹
 O. Hashimoto,¹ T. Hayakawa,² E. Hiyama,⁵ H. Hotchi,^{9,†} K. Imai,³ W. Imoto,⁹
 Y. Kakiguchi,⁵ M. Kameoka,¹ T. Kishimoto,² A. Krutenkova,¹⁰ T. Maruta,⁵
 A. Matsumura,¹ M. May,⁴ S. Minami,² Y. Miura,¹ K. Miwa,³ T. Miyoshi,^{1,‡}
 K. Mizunuma,¹ T. Nagae,⁵ S.N. Nakamura,¹ K. Nakazawa,¹¹ M. Niiyama,³
 H. Nomura,¹ H. Noumi,⁵ Y. Okayasu,¹ S. Ota,³ T. Ohtaki,⁹ H. Outa,^{5,§} P. Pile,⁴
 B.P. Quinn,⁷ A. Rusek,⁴ P.K. Saha,⁹ Y. Sato,⁵ T. Saitoh,⁶ M. Sekimoto,⁵
 R. Sutter,⁴ H. Takahashi,^{3,¶} T. Takahashi,¹ L. Tang,¹² K. Tanida,¹³ S. Terashima,³
 M. Togawa,³ A. Toyoda,⁵ M. Ukai,¹ H. Yamauchi,¹ L. Yuan,¹² S.H. Zhou¹⁴

¹ Department of Physics, Tohoku University, Sendai 980-8578, Japan

² Department of Physics, Osaka University, Toyonaka 560-0043, Japan

³ Department of Physics, Kyoto University, Kyoto 606-8502, Japan

⁴ Brookhaven National Laboratory, NY 11973, USA

⁵ Institute of Particle and Nuclear Studies, KEK, Tsukuba 305-0801, Japan

⁶ GSI, Darmstadt D-64291, Germany

⁷ Carnegie Mellon University, Pittsburgh, PA 15213, USA

⁸ Department of Physics, University of Freiburg, Freiburg 79104, Germany

⁹ Osaka Electro-Communication University, Neyagawa, 572-8530 Japan

¹⁰ Institute for Theoretical and Experimental Physics, Moscow, 117218 Russia

¹¹ Department of Physics, Gifu University, Gifu 501-1193, Japan

¹² Department of Physics, Hampton University, Hampton, VA 23668, USA

¹³ RIKEN, Wako 351-0198, Japan

¹⁴ China Institute of Atomic Energy, Beijing 102413, China
 (E930('01), E509, E518 collaborations)

The present status of hypernuclear γ -ray spectroscopy with Hyperball is summarized. We observed two γ transitions of $^{16}_{\Lambda}\text{O}(1^- \rightarrow 1^-, 0^-)$ and obtained the strength of the ΛN tensor force. In $^{10}\text{B}(K^-, \pi^-\gamma)$ data, we did not observe the spin-flip M1 transition of $^{10}_{\Lambda}\text{B}(2^- \rightarrow 1^-)$, but γ rays from hyperfragments such as $^7_{\Lambda}\text{Li}(7/2^+ \rightarrow 5/2^+)$ and $^9_{\Lambda}\text{Be}(3/2^+ \rightarrow 1/2^+)$ were observed. In $^{11}\text{B}(\pi^+, K^+\gamma)$ data, we observed six γ transitions of $^{11}_{\Lambda}\text{B}$. We also attempted an inclusive γ -ray measurement with stopped K^- beam.

[†]Present address: Japan Atomic Energy Research Institute, Tokai 319-1195, Japan.

[‡]Present address: University of Houston, Houston, TX 77204-5506, USA.

[§]Present address: RIKEN, Wako 351-0198, Japan.

[¶]Present address: KEK, Tsukuba 305-0801, Japan.

*Supported by Grant-In-Aid for Scientific Research from Ministry of Education of Japan, No.11440070 and 15204014.

1. Introduction

Since 1998, our project of hypernuclear γ spectroscopy with a germanium (Ge) detector array, Hyperball, has brought great progress in hypernuclear physics by revealing precise structure of several light Λ hypernuclei with a resolution in the keV range.

Hyperball is a large-acceptance germanium (Ge) detector array dedicated to hypernuclear γ spectroscopy. It has a large efficiency of 2.5 % at 1 MeV realized with fourteen large-volume Ge detectors, and is featured by special readout electronics which enables detection of γ rays under extremely high counting-rate conditions in hypernuclear experiments with meson beams. Details of Hyperball are described in Ref. [1,2].

One of the the most important physics motivations of hypernuclear γ spectroscopy is the study the ΛN interaction. In particular, we have been investigating the strengths of the ΛN spin-dependent forces from precise level structure of p -shell Λ hypernuclei. The potential of the ΛN two-body effective interaction can be written as:

$$V_{\Lambda N}^{eff}(r) = V_0(r) + V_\sigma(r)\boldsymbol{\sigma}_\Lambda\boldsymbol{\sigma}_N + V_\Lambda(r)\mathbf{l}_{\Lambda N}\boldsymbol{\sigma}_\Lambda + V_N(r)\mathbf{l}_{\Lambda N}\boldsymbol{\sigma}_N + V_T(r) [3(\boldsymbol{\sigma}_\Lambda\hat{\mathbf{r}})(\boldsymbol{\sigma}_N\hat{\mathbf{r}}) - \boldsymbol{\sigma}_\Lambda\boldsymbol{\sigma}_N]$$

The four spin-dependent terms, namely, the spin-spin term V_σ , the Λ -spin-dependent spin-orbit term V_Λ , the nucleon-spin-dependent spin-orbit term V_N , and the tensor term V_T , have not been studied well by experiments of the (K^-, π^-) and (π^+, K^+) reaction spectroscopy. The radial integrals of V_σ , V_Λ , V_N , and V_T with the $p_N s_\Lambda$ wavefunction in p -shell hypernuclei are denoted as Δ , S_Λ , S_N , and T , respectively. These effective-interaction parameters can be experimentally determined from low-lying level energies of p -shell hypernuclei [3,4]. However, because of a small level spacing between the spin-doublet states, high-resolution γ -ray spectroscopy with Ge detectors is necessary to investigate them.

1.1. Recent Experiments

Table 1 shows all the Hyperball experiments we have carried out. In 1998, we performed two experiments, KEK E419 for ${}^7_\Lambda\text{Li}$ [1,5,6] and BNL E930('98) for ${}^9_\Lambda\text{Be}$ [7], as we already reported in the HYP2000 conference. After that, we carried out the second run of BNL E930 (E930('01)) for ${}^{16}\text{O}$ and ${}^{10}\text{B}$ targets. With the ${}^{16}\text{O}$ target data, we observed ${}^{16}_\Lambda\text{O}$ and ${}^{15}_\Lambda\text{N}$ γ rays as described later. With the ${}^{10}\text{B}$ target data, we did not observe the ${}^{10}_\Lambda\text{B}(2^- \rightarrow 1^-)$ γ ray transition, but observed several γ rays from hyperfragments such

Table 1
List of experiments for hypernuclear γ spectroscopy with Hyperball since 1998.

Experiment	Year	Line	Target/Reaction	Hypernuclei studied
KEK E419	1998	K6	${}^7\text{Li}(\pi^+, K^+\gamma)$	${}^7_\Lambda\text{Li}$ [1,5,6]
BNL E930('98)	1998	D6	${}^9\text{Be}(K^-, \pi^-\gamma)$	${}^9_\Lambda\text{Be}$ [7]
BNL E930('01)	2001	D6	${}^{16}\text{O}(K^-, \pi^-\gamma)$ ${}^{10}\text{B}(K^-, \pi^-\gamma)$	${}^{16}_\Lambda\text{O}$, ${}^{15}_\Lambda\text{N}$ ${}^{10}_\Lambda\text{B}$, ${}^9_\Lambda\text{Be}$, ${}^7_\Lambda\text{Li}$ etc. [8]
KEK E509	2002	K5	${}^7\text{Li}, {}^9\text{Be}, {}^{10}\text{B}, {}^{11}\text{B}, {}^{12}\text{C}(K_{stop}^-, \gamma)$	hyperfragments (${}^7_\Lambda\text{Li}$) [9]
KEK E518	2002	K6	${}^{11}\text{B}(\pi^+, K^+\gamma)$	${}^{11}_\Lambda\text{B}$ [10]




Figure 1. Level schemes of ${}^7_\Lambda\text{Li}$, ${}^9_\Lambda\text{Be}$, ${}^{10}_\Lambda\text{B}$, ${}^{11}_\Lambda\text{B}$, and ${}^{16}_\Lambda\text{O}$ determined from Hyperball experiments. Newly observed γ rays, measured level energies, and assigned spins in the recent experiments (E930('01), E509, E518) are shown in thick arrows and bold letters.

as ${}^7_\Lambda\text{Li}$ and ${}^9_\Lambda\text{Be}$. In 2002, we moved Hyperball from BNL to KEK and performed two experiments, E509 for hyperfragments [9] and E518 for ${}^{11}_\Lambda\text{B}$ [10].

Figure 1 shows the level schemes of p -shell hypernuclei determined from these Hyperball experiments. The γ rays first observed and identified in E930('01), E509, and E518, which are shown in thick arrows, are described in detail in the following sections.

2. ${}^9_\Lambda\text{Be}$ and the spin-orbit force

The experiment BNL E930 aims at determination of all the spin-dependent force strengths from structure of several p -shell hypernuclei. Using high-intensity and pure K^- beam at 0.93 GeV/c provided by the D6 beam line at BNL AGS, hypernuclei were produced by the (K^-, π^-) reaction. The momenta of incident K^- and scattered π^- were measured with magnetic spectrometers to obtain the hypernuclear excitation spectrum. γ rays were detected with Hyperball installed around the target.

We previously reported that the ${}^9_\Lambda\text{Be}$ target data in E930('98) exhibited the $5/2^+, 3/2^+ \rightarrow 1/2^+$ transitions and revealed a hypernuclear fine structure of ${}^9_\Lambda\text{Be}(5/2^+, 3/2^+)$ [7]. Recently, we have applied Doppler-shift correction to this ${}^9_\Lambda\text{Be}$ spectrum and observed clearly-




Figure 2. Doppler-shift corrected spectrum of ${}^9_{\Lambda}\text{Be}$ γ rays around 3 MeV obtained in the E930('98) experiment. The two-peak structure was well fitted by the simulated peak shape with a lifetime of < 0.1 ps.




Figure 3. γ -ray spectrum for the mass region slightly higher than the bound-state region of ${}^{10}_{\Lambda}\text{B}$ ($-18 < -B_{\Lambda} < 28$ MeV). A γ -ray peak from ${}^9_{\Lambda}\text{Be}$ is observed.

separated two peaks as shown in Fig. 2. This structure was well fitted by the simulated peak shape with a short lifetime (< 0.1 ps). The γ -ray energies were obtained to be $3024 \pm 3 \pm 1$ and $3067 \pm 3 \pm 1$ keV, and the separation energy to be 43 ± 5 keV. The separation energy and the lifetime have been revised from the previous values in Ref. [7].

In the ${}^{10}\text{B}$ target data taken in E930('01), we observed a γ -ray peak at 3065 keV, when the mass region slightly higher than the ${}^{10}_{\Lambda}\text{B}$ bound states is selected, as shown in Fig. 3. ${}^9_{\Lambda}\text{Be}$ can be produced by proton emission from excited states of ${}^{10}_{\Lambda}\text{B}$, and this observed γ -ray energy coincides with the energy of one of the ${}^9_{\Lambda}\text{Be}$ γ rays observed in the ${}^9\text{Be}$ target run. Therefore, this γ ray is assigned as one of the ${}^9_{\Lambda}\text{Be}$ transitions of $3/2^+ \rightarrow 1/2^+$ or $5/2^+ \rightarrow 1/2^+$. The ${}^{10}_{\Lambda}\text{B}(3^-, -B_{\Lambda} \sim 1 \text{ MeV})$ state, which is expected to have a large cross section, mostly decays into ${}^9_{\Lambda}\text{Be}(3/2^+) + p$ (see Fig. 1 (d)), while other ${}^{10}_{\Lambda}\text{B}$ excited states decaying into ${}^9_{\Lambda}\text{Be}(5/2^+) + p$ have much smaller cross sections [11]. Therefore, we assigned the observed peak as $3/2^+ \rightarrow 1/2^+$. From this spin assignment, the previous result for the Λ -spin-dependent spin-orbit force parameter of $-0.02 < S_{\Lambda} < 0.03$ MeV [7] was improved to $-0.02 < S_{\Lambda} < -0.01$ MeV. This sign of the Λ -spin-dependent spin-orbit term is consistent with the $p_{1/2} - p_{3/2}$ spin-orbit splitting in ${}^{13}_{\Lambda}\text{C}$ measured with NaI counter arrays by the BNL E929 experiment [12].

3. $^{16}_{\Lambda}\text{O}$ and ΛN tensor force

The purpose of the ^{16}O target run in E930('01) is to investigate the ΛN tensor force strength (T), which has never been studied experimentally. Since the one-pion exchange is forbidden in the ΛN interaction, the tensor force is expected to be small, but the kaon exchange and the two-pion exchange through the Σ - Λ coupling are expected to give some contribution to the tensor force.

It was pointed out that energy spacings of the spin doublets in $p_{1/2}$ -shell hypernuclei are sensitive to the ΛN tensor force strength [3]. According to a shell-model calculation by Millener, the spacing of the ground-state doublet ($0^-, 1^-$) of $^{16}_{\Lambda}\text{O}$ is given as [11]

$$E(1^-) - E(0^-) = -0.382\Delta + 1.378S_{\Lambda} - 0.004S_N + 7.850T + \Lambda\Sigma \text{ (MeV)}, \quad (1)$$

where $\Lambda\Sigma$ denotes the effect of $\Lambda - \Sigma$ coupling. By the $^{16}\text{O}(K^-, \pi^-)$ reaction, we can populate the 6 MeV-excited $^{16}_{\Lambda}\text{O} [(p_{3/2})_n^{-1}(s_{1/2})_{\Lambda}]_{1^-}$ state and detect $M1$ transitions from this state to each member of the ground-state doublet, even if the spacing is too small (< 100 keV) to detect the spin-flip $M1$ transition between the doublet members (see Fig. 1 (c)). In addition, the 11 MeV-excited $[(p_{1/2})_n^{-1}(p_{1/2})_{\Lambda}]_{0^+}$ state and the 17 MeV-excited $[(p_{3/2})_n^{-1}(p_{3/2})_{\Lambda}]_{0^+}$ state of $^{16}_{\Lambda}\text{O}$ are expected to decay to excited states of $^{15}_{\Lambda}\text{N}$ by proton emission with sizable branching ratios, which is followed by emission of $^{15}_{\Lambda}\text{N}$ γ rays. The ground-state doublet spacing of $^{15}_{\Lambda}\text{N}$, which also has a large contribution of the ΛN tensor force, may also be measured.

The experimental method and setup are almost identical to those in the previous E930 run for $^9_{\Lambda}\text{Be}$ described in Ref. [7]. We used a 20 cm-thick water target and irradiated it with 4.0×10^{10} K^- in total. More description on this experiment is found in Ref. [8].

Figure 4 shows preliminary γ -ray spectra for $^{16}_{\Lambda}\text{O}$. Figure 4 (a) shows the spectrum when events in the 6 MeV-excited 1^- state region of the $^{16}_{\Lambda}\text{O}$ mass spectrum ($-21 < -B_{\Lambda} < 8$ MeV) are selected. A broad bump is observed at around 6.55 MeV. After the event-by-event Doppler-shift correction was applied, the broad bump is resolved into two narrower peaks as shown in (b). This structure is not observed for the highly unbound region ($-B_{\Lambda} > 50$ MeV) as shown in Fig. 4 (c), in which beam-induced γ rays from the ^{16}O target are observed. (The 6130 keV ^{16}O peak width demonstrates the resolution in this energy region.)

The structure at 6.55 MeV is thus attributed to the $M1(1_2^- \rightarrow 1_1^-, 0^-)$ transitions in $^{16}_{\Lambda}\text{O}$. The peaks in Fig. 4 (b) were fitted with the expected Doppler-corrected peak shape which was calculated from a simulation for the Doppler-shift correction. The spectrum was fitted well with two peaks as shown in Fig. 5. The energies (and the counts) of these peaks were obtained as 6534.1 ± 1.5 keV (149 ± 18 counts) and 6560.2 ± 1.3 keV (226 ± 30 counts). By comparing the ratio of the peak counts with the expected branching ratios, the 6534 keV and 6560 keV peaks were assigned as $1_2^- \rightarrow 1_1^-$ and $1_2^- \rightarrow 0^-$ transitions, respectively [8]. Then we obtained the energy spacing of the ground-state doublet:

$$E(1^-) - E(0^-) = 26.1 \pm 2.0 \text{ keV (preliminary)}.$$

It is the smallest spacing in hypernuclear fine structure observed so far. This very small spacing results from a cancellation of the spin-spin force (Δ term) and the tensor force (T term) contributions. It gives the tensor term strength of $T = +30$ keV (preliminary) from




Figure 4. γ -ray spectrum of $^{16}_\Lambda\text{O}$ (preliminary). (a) Bound-state region ($-21 < -B_\Lambda < 8$ MeV) is gated. (b) Same as (a) but Doppler-shift correction is applied. (c) Highly unbound region ($-B_\Lambda > 50$ MeV) is gated.




Figure 5. (a) Simulated peak shape for a fast γ transition after Doppler-shift correction. (b) The structure around 6.55 MeV in Fig. 4 (b) was fitted with two peaks of the simulated peak shape (preliminary).

Eq. 1 and with the Δ , S_Λ , and S_N values already determined from previous Hyperball experiments. This is the first experimental information on the ΛN tensor force.

The meson-exchange baryon-baryon interactions models (ND, NF, NSC89, NSC97f) predict a tensor force strength of $T = 18 - 54$ keV through a G-matrix calculation [13]. They are almost consistent with the experimental value.

4. Study of $^{10}_\Lambda\text{B}$

The purpose of the ^{10}B target run in E930('01) is to measure the energy spacing of the $^{10}_\Lambda\text{B}$ ground-state doublet ($2^-, 1^-$) by observing the spin-flip $M1$ transition ($2^- \rightarrow 1^-$). Since the production cross section of the 2^- state is large enough, the $M1$ transition can be easily observed if the level spacing is as large as predicted (~ 200 keV). On the other hand, if the spacing is smaller than ~ 100 keV, the γ transition is overcome by weak decay.

Figure 6 is a preliminary γ -ray spectrum when the bound-state region of $^{10}_\Lambda\text{B}(K^-, \pi^-)_\Lambda\text{B}$ is selected. We observed no peak structure. Considering the number of the expected γ -ray peak yield, we concluded that the 2^- state is higher than the 1^- state only by 100 keV or less, or the order of the spins in the doublet is reversed. Thus, we confirmed the old result by Chrien *et al.* [14] with higher statistics. The confirmed result of $E(2^-) - E(1^-) < 100$ keV seems contradictory to the Δ value (0.4 MeV) obtained from the $^7_\Lambda\text{Li}$ ground-state




Figure 6. Preliminary γ -ray spectrum for the bound-state region of $^{10}_{\Lambda}\text{B}$ ($-23 < -B_{\Lambda} < 7$ MeV). No peak is observed.

doublet ($3/2^+, 1/2^+$) spacing, suggesting that more theoretical and experimental studies are necessary, particularly for the Σ - Λ coupling effect as investigated in Ref. [11].

5. Complete study of $^7_{\Lambda}\text{Li}$

In the ^{10}B target data in E930('01), we observed γ rays from $^7_{\Lambda}\text{Li}$ produced as hyperfragments from highly excited states of $^{10}_{\Lambda}\text{B}$, presumably through the s -substitutional ^{10}B (3^+ , ~ 28 MeV excited) state decaying into $^7_{\Lambda}\text{Li} + ^3\text{He}$ as shown in Fig. 1 (d). Figure 7 (a) shows the γ -ray spectrum when the unbound region ($0 < -B_{\Lambda} < 40$ MeV) was selected. The $M1(3/2^+ \rightarrow 1/2^+)$ and $E2(5/2^+ \rightarrow 1/2^+)$ γ rays of $^7_{\Lambda}\text{Li}$ previously observed in E419 are identified. We selected the $E2$ γ -ray events ($2042 < E_{\gamma} < 2058$ keV) and plotted a spectrum of another γ ray emitted in coincidence. As shown in Fig. 7 (b), a peak was observed at 471 keV. The probability that background fluctuation makes such a peak anywhere in the region of 0.1–1 MeV is 0.006%. This peak is assigned as the $M1(7/2^+ \rightarrow 5/2^+)$ transition, because it is the only transition emitted in coincidence with $E2(5/2^+ \rightarrow 1/2^+)$. This is the first successful application of the γ - γ coincidence method to hypernuclei.

The observed energy can be compared with theoretical predictions. A cluster-model calculation by Hiyama *et al.* predicted the ($7/2^+, 5/2^+$) spacing to be 560 keV when the Λ -spin-dependent spin-orbit force is assumed to be zero [15]. It is close to the observed value. According to the Millener's shell-model calculation [11], the energy spacing is described as $E(7/2^+) - E(5/2^+) = 1.29\Delta + 2.20S_{\Lambda} + 0.02S_N - 2.39T + \Lambda\Sigma$. By using the already determined values of Δ , S_{Λ} , S_N , T , and the theoretically calculated $\Lambda\Sigma$ value, the equation gives 511 keV, being also close to the observation. It is found that the observed value is consistent with the already-known strengths of the spin-spin force (Δ) and the very small spin-orbit force (S_{Λ}).

Together with the E419 results [1], we have clarified the complete level scheme and energies of all the bound states of $^7_{\Lambda}\text{Li}$ as shown in Fig. 1 (a).

6. Spectroscopy of $^{11}_{\Lambda}\text{B}$ (E518)

In 2002, we carried out a γ spectroscopy experiment of $^{11}_{\Lambda}\text{B}$ with the (π^+, K^+) reaction at 1.05 GeV/c employing Hyperball and the SKS spectrometer at KEK-PS [10]. One of the purposes of this experiment is to measure the transition probability $B(M1)$ of




Figure 7. (a) γ -ray spectrum for the unbound region ($0 < -B_\Lambda < -40$ MeV) of $^{10}_\Lambda\text{B}$. Two γ -ray peaks from $^7_\Lambda\text{Li}$ produced as hyperfragments are observed. (b) γ -ray spectrum in coincidence with the $^7_\Lambda\text{Li}$ $E2(5/2^+ \rightarrow 1/2^+)$ γ -ray peak.

the Λ spin-flip $M1$ transition $^{11}_\Lambda\text{B}(3/2^+ \rightarrow 1/2^+)$ and extract information on the magnetic moment of a Λ inside a nucleus by the method described in Ref. [16]. The other purpose is to cross-check the ΛN spin-dependent interaction parameters which have been determined from the $^7_\Lambda\text{Li}$, $^9_\Lambda\text{Be}$, and $^{16}_\Lambda\text{O}$ experiments with Hyperball.

The experimental setup is almost identical to the one in E419 [1,5]. We used a 10 cm-thick 98%-enriched ^{11}B metal target. When the bound-state region is gated in the $^{11}_\Lambda\text{B}$ mass spectrum, the γ -ray spectrum exhibited six peaks as shown in Fig. 8. One of them was observed in the Doppler-shift-corrected spectrum. They are attributed to transitions from $^{11}_\Lambda\text{B}$, but the assignment of all these γ rays and the reconstruction of the level scheme are difficult because of low statistics which does not allow $\gamma\gamma$ coincidence measurement.

The prominent peak at 1482 keV is assigned as $E2(1/2^+ \rightarrow 5/2^+)$ (see Fig. 1 (e)). It is likely an $E2$ transition because its narrow width indicates a lifetime of the transition longer than ~ 10 ps, which gives a very small $B(M1)$ value if it is an $M1$ transition. The $1/2^+ \rightarrow 5/2^+$ transition is the only $E2$ transition expected in $^{11}_\Lambda\text{B}$, and the observed largest γ -ray yield is also consistent with this assignment. It is to be noted that the shell-model prediction by Millener [11] for this $E2$ energy with the experimentally determined ΛN interaction parameters is 1020 keV, significantly lower than the observed energy.

7. Hyperfragments (E509)

In KEK-PS E509, we attempted an experiment of inclusive γ -ray measurement in the stopped K^- absorption reaction, which is known to produce various hyperfragments with large production yields. See Ref. [9] for details. We stopped K^- from the K5 beam line on several light targets (^7Li , ^9Be , ^{10}B , ^{11}B , and ^{12}C) and measured γ rays with Hyperball.




Figure 8. Preliminary γ ray spectra of $^{11}\text{B}(\pi^+, K^+\gamma)^{11}_{\Lambda}\text{B}$ experiment at KEK (E518). Six γ rays of $^{11}_{\Lambda}\text{B}$ are observed.




Figure 9. γ ray spectra in the (K_{stop}^-, γ) reaction on ^7Li and ^{10}B targets. The 2050 keV γ ray from $^7_{\Lambda}\text{Li}$ is abundantly observed in the ^{10}B target spectrum.

From the ^{10}B , ^{11}B , and ^{12}C targets, we observed the $^7_{\Lambda}\text{Li}(5/2^+ \rightarrow 1/2^+)$ transition at 2050 keV. The yield of this γ ray for the ^{10}B target is very large, 500 counts in 3.5 days' beam time, suggesting the effectiveness of this method. The production rate of the $^7_{\Lambda}\text{Li}(5/2^+)$ state is derived to be $0.075 \pm 0.016\%$ per stopped K^- on ^{10}B target.

This method is powerful with a larger Ge detector array, by which the $\gamma\gamma$ coincidence method allows detection of hypernuclear γ rays and their assignments. It may open a new possibility to study various hypernuclei including neutron/proton-rich ones which cannot be produced by the direct reactions such as (K^-, π^-) and (π^+, K^+) .

8. Future Plans

We are now constructing Hyperball2, an upgraded version of Hyperball for near-future experiments at KEK and BNL. It has an efficiency twice as large as the present Hyperball,

realizing a $\gamma\gamma$ coincidence efficiency larger by four times. It will play an essential role in study of hypernuclei having complicated level schemes such as ${}^{\Lambda}_{11}\text{B}$.

At the 50 GeV proton accelerator facility at J-PARC, we plan to pursue various types of hypernuclear γ spectroscopy experiments [17,18]. We have started development of faster readout techniques and faster background-suppression counters necessary for the stronger beams expected at J-PARC.

9. Summary

We have investigated various p -shell Λ hypernuclei employing Hyperball. In E930('01), we successfully observed two γ transitions of $1_2^- \rightarrow 1_1^-, 0^-$ at 6.55 MeV, and the ground-state doublet ($1_1^-, 0^-$) spacing was obtained to be $E(1_1^-) - E(0^-) = 26.1 \pm 2.0$ keV. It gives the ΛN tensor force strength of $T = +30$ keV. All the ΛN spin-dependent force parameters have been thus determined from our γ spectroscopy experiments. In ${}^{10}\text{B}(K^-, \pi^-\gamma)$ data, we observed γ rays from hyperfragments such as ${}^9_{\Lambda}\text{Be}(3/2^+ \rightarrow 1/2^+)$ and ${}^7_{\Lambda}\text{Li}(7/2^+ \rightarrow 5/2^+)$. In the observation of the ${}^7_{\Lambda}\text{Li}(7/2^+ \rightarrow 5/2^+)$ transition, we successfully applied the $\gamma\gamma$ coincidence method to hypernuclei for the first time. On the other hand, the spin-flip M1 transition of ${}^{10}_{\Lambda}\text{B}(2^- \rightarrow 1^-)$ was not observed. At KEK, we studied ${}^{11}_{\Lambda}\text{B}$ with the (π^+, K^+) reaction and observed six γ transitions. We also performed a pioneering experiment with the $(K_{\text{stop}}^-, \gamma)$ reaction (E509), and observed a γ -ray peak from ${}^7_{\Lambda}\text{Li}$ hyperfragments. Hypernuclear γ spectroscopy will be further pursued with much stronger beams at J-PARC.

REFERENCES

1. H. Tamura *et al.*, Phys. Rev. Lett. 84 (2000) 5963.
2. K. Tanida, Ph.D thesis, University of Tokyo (2000).
3. R.H. Dalitz and A. Gal, Ann. Phys. 116 (1978) 167.
4. D.J. Millener, A. Gal, C.B. Dover and R.H. Dalitz, Phys. Rev. C 31 (1985) 499.
5. K. Tanida *et al.*, Phys. Rev. Lett. 86 (2001) 1982.
6. J. Sasao *et al.*, Phys. Lett. B 579 (2004) 258.
7. H. Akikawa *et al.*, Phys. Rev. Lett. 88 (2002) 082501.
8. M. Ukai *et al.*, in these proceedings.
9. K. Tanida *et al.*, Nucl. Phys. A 721 (2003) 999c; K. Miwa *et al.*, in these proceedings.
10. Y. Miura *et al.*, in these proceedings.
11. D.J. Millener, in these proceedings.
12. S. Ajimura *et al.*, Phys. Rev. Lett. 86 (2001) 4255.
13. D.J. Millener, Proc. Jlab Sponsored Workshop on *Hypernuclear Physics with Electromagnetic Probes*, Ed. L. Tang and O. Hashimoto, Hampton, December 1999, p.79.
14. R.E. Chrien *et al.*, Phys. Rev. C 41 (1990) 1062.
15. E. Hiyama *et al.*, Nucl. Phys. A 639 (1998) 173c.
16. H. Tamura, Nucl. Phys. A 691 (2001) 86c.
17. K. Imai *et al.*, Letter of Intent for Nuclear and Particle Physics Experiments at the J-PARC, KEK, L06 (2003), <http://www-ps.kek.jp/jhf-np/L0Ilist/L0Ilist.html>
18. T. Nagae, in these proceedings.